

Cav1.2 Antibody
Cav1.2 Antibody, Clone S57-46
Catalog # ASM10178**Specification**

Cav1.2 Antibody - Product Information

Application	IHC, IP, WB
Primary Accession	P15381
Other Accession	NP_001129994.1
Host	Mouse
Isotype	IgG1
Reactivity	Human, Mouse, Rat, Hamster
Clonality	Monoclonal

Description

Mouse Anti-Rabbit Cav1.2 Monoclonal IgG1

Target/Specificity

Detects ~240kDa (varies with cell background due to glycosylation).

Other Names

CACH3 Antibody, CACN4 Antibody, CACNA 1D Antibody, CACNL1A2 Antibody, voltage dependent L type calcium channel subunit alpha 1D Antibody, alpha-1 subunit voltage-dependent calcium channel Antibody, calcium channel voltage-dependent L type alpha 1C subunit1 Antibody, voltage-gated calcium channel alpha subunit Cav1.2 Antibody, calcium channel L type Antibody, alpha 1 polypeptide isoform 1 cardiac muscle Antibody, calcium channel cardiac dihydropyridine-sensitive alpha-1 subunit Antibody, voltage-gated L-type calcium channel Cav1.2 alpha 1 subunit splice variant 10 Antibody, DHPR alpha-1 subunit Antibody, Voltage-gated calcium channel subunit alpha Cav1.2 Antibody, Calcium channel L type alpha-1 polypeptide isoform 1 cardiac muscle Antibody

Immunogen

Fusion protein amino acids 1507-1733 (intracellular carboxyl terminus) of rabbit Cav1.2

Purification

Protein G Purified

Storage **-20°C****Storage Buffer**

PBS pH7.4, 50% glycerol, 0.1% sodium azide

Shipping Temperature **Blue Ice or 4°C****Certificate of Analysis**

1 µg/ml of SMC-300 was sufficient for detection of Cav1.2 in 10 µg of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

Cellular Localization


Membrane | Cell Membrane

Cav1.2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cav1.2 Antibody - Images




Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Cav1.2 Monoclonal Antibody, Clone S57-46 (ASM10178). Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-Cav1.2 Monoclonal Antibody (ASM10178) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:200 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min at RT, 5 min at RT. Localization: Cell Membrane, Membrane, Cytoplasm, Nucleoplasm. Magnification: 60X. (A) Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain. (B) Anti-Cav1.2 Antibody. (C) Composite. (D) Composite.




Immunohistochemistry analysis using Mouse Anti-Cav1.2 Calcium Channel Monoclonal Antibody, Clone S57-46 (ASM10178). Tissue: hippocampus. Species: Human. Fixation: 10% formalin. Primary Antibody: Mouse Anti-Cav1.2 Calcium Channel Monoclonal Antibody (ASM10178) at 1:100 for 1


hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.



Immunoprecipitation analysis using Mouse Anti-CaV1.2 Calcium Channel Monoclonal Antibody, Clone S57-46 (ASM10178). Tissue: INS-1E cells. Species: Rat. Primary Antibody: Mouse Anti-CaV1.2 Calcium Channel Monoclonal Antibody (ASM10178) at 1:200. Courtesy of: Merrie Mosedale.



Western Blot analysis of Hamster T-CHO cell lysate showing detection of CaV1.2 Calcium Channel protein using Mouse Anti-CaV1.2 Calcium Channel Monoclonal Antibody, Clone S57-46 (ASM10178). Primary Antibody: Mouse Anti-CaV1.2 Calcium Channel Monoclonal Antibody (ASM10178) at 1:1000.



Immunohistochemistry analysis using Mouse Anti-CaV1.2 Calcium channel Monoclonal Antibody, Clone S57-47 (ASM10178). Tissue: Brain Tissue. Species: Mouse. Fixation: Formalin. Primary Antibody: Mouse Anti-CaV1.2 Calcium channel Monoclonal Antibody (ASM10178) at 1:10000 for 12 hours at 4°C. Secondary Antibody: Biotin Goat Anti-Mouse at 1:2000 for 1 hour at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 200 µl for 2 minutes at RT. Magnification: 40x.

Cav1.2 Antibody - Background

Cav1.2 is a cardiac L-type calcium channel, and is important for excitation and contraction of the heart (1). It may be associated with a variant of Long QT syndrome called Timothy's syndrome (2, 3) and also with Brugada syndrome. Some references also suggest it is related to bipolar disease as well (3).

Cav1.2 Antibody - References

1. Splawski I., et al. (2004) Cell. 119 (1): 19-31.
2. Krey J.F., and Dolmetsch R. (2009) Biophysical. 96 (3): 221a-222a.
3. Crotti L., Celano G., Dagradi F. and Schwartz P.J. (2008) Orphanet J Rare Disease 3:18.