

**Monoclonal Antibody to Human PD-L1 (Clone : ABM4E54)**

#Cat: NB-30-00125    Size: 100ul

**Clonality:** Monoclonal**Clone Name:** ABM4E54**Application:** IHC, FACS, WB**Reactivity:** Human**Gene:** CD274**Gene ID:** 29126**Uniprot ID:** Q9NZQ7**Format:** Purified**Alternative Name:** CD274, B7H1, PDCD1L1, PDCD1LG1, PDL1**Isotype:** Mouse IgG2a Kappa**Immunogen Information :** A partial length recombinant PDL1 protein (amino acids 18-227) was used as the immunogen for this antibody.**Description:**

PD-L1 (CD274/B7-H1) is a critical membrane-bound costimulatory molecule belonging to the B7 superfamily that inhibits immune responses through its receptor, PD-1. PD-L1 plays a key role in the pathogenesis of inflammatory diseases (programmed death 1). It is widely expressed in the mononuclear phagocyte system (MPS), may co-stimulate T cells, and regulates inflammatory responses. PD-L1 exerts inflammation regulatory functions via a negative co-stimulatory effect on T cell functions to inhibit cytokine secretion, facilitate apoptosis of activated T cells and induce T cell anergy. Aberrant expression and dysregulation of CD274/PD-L1 have been reported during bacterial infection, inflammation, and in numerous autoimmune diseases.

**Product Info:****Amount:** 25 µg / 100 µg**Purification:** Protein G Chromatography**Content:** 25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.**Storage condition:** Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.**Application Note:**

Western blot analysis: 2-4 µg/ml; Immunohistochemical analysis-5-10 µg/ml; FACS: 1-2 µg/ml




Figure-1: Cell Surface FLOW analysis of PD-L1 in PHA treated human PBMC using 1  $\mu\text{g}$  of PD-L1 antibody (Clone: ABM4E54). Green represents isotype control; red represents anti-PD-L1 antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Figure-2: Immunohistochemical analysis of PD-L1 in Hodgkin's Lymphoma tissue using PD-L1 antibody (Clone: ABM4E54) at 5  $\mu\text{g}/\text{ml}$ .




Figure-3: Western blot analysis of PD-L1. Anti-PD-L1 antibody (Clone: ABM4E54) was tested at 0.5  $\mu\text{g}/\text{ml}$  on Recombinat lysates.




Figure-4: Western blot analysis of PD-L1. Anti-PD-L1 antibody (Clone: ABM4E54) was tested at 2  $\mu\text{g}/\text{ml}$  on (1) A549, (2) MCF-7, (3) 293, (4) HCT-116, (5) Saos2 and (6) Hela lysates.




Figure-5: Immunohistochemical analysis of PD-L1 in Human Tonsil tissue using PD-L1 antibody (Clone: ABM4E54) at 5  $\mu\text{g/ml}$ .




Figure-6: Western blot analysis of PD-L1. Anti-PD-L1 antibody (Clone: ABM4E54) was tested at 0.5  $\mu\text{g/ml}$  on (1) HepG2, (2) SKBR3, (3) A431, (4) THP1, (5) NCCIT, (6) PC3, (7) PANC-1, (8) U87 and (9) KATO-111 lysates.




Figure-7: Immunohistochemical analysis of PD-L1 in Human Lung Cancer tissue using PD-L1 antibody (Clone: ABM4E54).




Figure-8: Western blot analysis of PD-L1. Anti-PD-L1 antibody (Clone: ABM4E54) was tested at 2  $\mu\text{g/ml}$  on h Spleen lysate.