

Anti-Phospho-Ser⁷⁷⁴ Dynamin

Catalog Number: SY-p1050-774 Size: 100 μl

\$375.00


Product Description: Affinity purified sheep polyclonal antibody

Applications: WB: 1:1000 IF (fixed cultured neurons; Tan et al., 2003): 1:1000 IHC (fixed cultured neurons, Tan et al., 2003): 1:1000

Antigen: Phosphopeptide corresponding to amino acid residues surrounding the phospho-Ser⁷⁷⁴ of rat dynamin.

Specifies reactivity: The antibody has been directly tested for reactivity in Western blots with rat tissue. It is anticipated that the antibody will react with bovine, canine, chicken, human, mouse and non-human primates based on the fact that these species have 100% homology with the amino acid sequence used as antigen.

Biological Significance: Dynamin is a member of a group of nerve terminal proteins called dephosphins that regulate synaptic vesicle endocytosis (Cousin et al., 2001; Graham et al., 2002; Tsuboi et al., 2002). Cyclin dependent protein kinase 5 phosphorylates dynamin at Ser⁷⁷⁴ and Ser⁷⁷⁸ that are the phosphorylation sites on dynamin phosphorylated *in vivo* (Tan et al., 2003). Phosphorylation of these sites on dynamin is thought to play a key role in synaptic vesicle trafficking.



Western blot of rat hippocampal lysate stimulated with forskolin showing specific immunolabeling of the ~95k dynamin phosphorylated at Ser⁷⁷⁴ (Control). The phosphospecificity of this labeling is shown in the second lane (*lambda*-phosphatase: λ -Ptase). The blot is identical to the control except that it was incubated in λ -Ptase (1200 units for 30 min) before being exposed to the phospho-Ser⁷⁷⁴ dynamin. The immunolabeling is completely eliminated by treatment with λ -Ptase.

Page 1 of 2

WB = Western Blot **IF** = Immunofluorescence **IHC** = Immunohistochemistry **IP** = Immunoprecipitation **Packaging:** 100 μ I in 10 mM HEPES (pH 7.5), 150 mM NaCl, 100 μ g BSA per ml and 50% glycerol. Adequate amount of material to conduct 10-mini Western Blots.

Storage and Stability. For long term storage -20°C is recommended. Stable at -20°C for at least 1 year. **Shipment:** Domestic - Blue Ice; International – Dry Ice.

Purification Method: Prepared from sheep serum by affinity purification via sequential chromatography on phospho- and dephosphopeptide affinity columns.

Antibody Specificity: Specific for the ~95k dynamin protein phosphorylated at Ser⁷⁷⁴. Labels the purified protein phosphorylated *in vitro* by cdk5 but not by PKC. Does not cross react with other purified substrates of cdk5 (*e.g.* amphiphysin and synapsin). Immunolabeling is blocked by λ -phosphatase treatment.

Quality Control Tests: Western blots performed on each lot.

References:

- Cousin MA, Tan TC, Robinson PJ (2001) Protein phosphorylation is required for endocytosis in nerve terminals: potential role for the dephosphins dynamin I and synaptojanin, but not AP180 or amphiphysin. J Neurochem 76:105-116.
- Graham ME, O'Callaghan DW, McMahon HT, Burgoyne RD (2002) Dynamin-dependent and dynamin-independent processes contribute to the regulation of single vesicle release kinetics and quantal size. Proc Natl Acad Sci USA 99:7124-7129.
- Tan TC, Valova VA, Malladi CS, Graham ME, Berven LA, Jupp OJ, Hansra G, McClure SJ, Sarcevic B, Boadle RA, Larsen MR, Cousin MA, Robinson PJ (2003) cdk5 is essential for synaptic vesicle endocytosis. Nat Cell Biol 5:701-710.
- Tsuboi T, Terakawa S, Scalettar BA, Fantus C, Roder J, Jeromin A (2002) Sweeping model of dynamin activity -Visualization of coupling between exocytosis and endocytosis under an evanescent wave microscope with green fluorescent proteins. J Biol Chem 277:15957-15961.

Page 2 of 2