ABclonal www.abclonal.com

ABflo® 488 Rabbit IgG isotype control

Catalog No.: A22069

Basic Information

Observed MW

Calculated MW

Category

SMab Recombinant Monoclonal Antibody

Applications

FC

Cross-Reactivity

CloneNo number

ARC5105-10-ABf488

Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

Background

The isotype of a primary antibody and the application it is being used in can result in background staining. Primary antibody background noise can be caused by binding to Fc receptors on target cells; by non-specific interactions with cellular proteins, carbohydrates, and lipids; or by cell autofluorescence. Isotype control antibodies can act as negative controls to help differentiate non-specific background signal from specific antibody signal because they have no relevant specificity to a target antigen. An isotype control antibody should have the same immunoglobulin type and be used at the same concentration as the test antibody.

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID

Swiss Prot

Immunogen

A synthesized peptide derived from rabbit IgG isotype control

Synonyms

Contact

•

www.abclonal.com

Product Information

Source Rabbit **Isotype**


IgG

PurificationAffinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.



Flow cytometry: 1X10^6 Human PBMC(negative control) were surface-stained with ABflo® 488 Rabbit anti-Human CD13/ANPEP mAb(A21945,5 µl/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line).Non-fluorescently stained Human PBMC were used as blank control (red line).



Flow cytometry: 1X10^6 Human PBMC(negative control) were surface-stained with ABflo® 488 Rabbit anti-Human CD48 mAb(A21945,5 µl/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line).Non-fluorescently stained Human PBMC wereused as blank control (red line).