

RP2 antibody

Product Information

| Catalog No.: | FNab07390 |
|---------------|--|
| Size: | 100µg |
| Form: | liquid |
| Purification: | Immunogen affinity purified |
| Purity: | \geq 95% as determined by SDS-PAGE |
| Host: | Rabbit |
| Clonality: | polyclonal |
| Clone ID: | None |
| IsoType: | IgG |
| Storage: | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.) |

Background

Acts as a GTPase-activating protein(GAP) involved in trafficking between the Golgi and the ciliary membrane. Involved in localization of proteins, such as NPHP3, to the cilium membrane by inducing hydrolysis of GTP ARL3, leading to the release of UNC119(or UNC119B). Acts as a GTPase-activating protein(GAP) for tubulin in concert with tubulin-specific chaperone C, but does not enhance tubulin heterodimerization. Acts as guanine nucleotide dissociation inhibitor towards ADP-ribosylation factor-like proteins.


Immunogen information

| Immunogen: | retinitis pigmentosa 2 |
|--------------|------------------------|
| Synonyms: | Protein XRP2 RP2 |
| Observed MW: | 40 kDa |
| Uniprot ID : | 075695 |

Application

| Reactivity: | Human, Mouse, Rat | | |
|---|--------------------|--|--|
| Tested Application: | ELISA, WB, IHC, FC | | |
| Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200 | | | |
| Image: | | | |





Immunohistochemistry of paraffin-embedded human liver cancer using FNab07390(RP2 antibody) at dilution of 1:50

Jurkat cells were subjected to SDS PAGE followed by western blot with FNab07390(RP2 antibody) at dilution of 1:500