

Neuroligin 4 Antibody

Neuroligin 4 Antibody, Clone S98-7 Catalog # ASM10301

Specification

Neuroligin 4 Antibody - Product Information

Application	IHC
Primary Accession	<u>B0F2B4</u>
Other Accession	<u>ABS19580</u>
Host	Mouse
Isotype	lgG1
Reactivity	Mouse
Clonality	Monoclonal
Description	
Mouse Anti-Mouse Neuroligin 4 Monoclonal IgG1	

Target/Specificity

Detects ~125-130kDa. Does not cross-react with other Neuroligins.

Other Names

ASPGX2 Antibody, AUTSX2 Antibody, HLNX Antibody, HNLX Antibody, KIAA1260 Antibody, MGC22376 Antibody, NLGN Antibody, NLGN4 Antibody, NLGN4X Antibody, neuroligin-4 X-linked Antibody, neuroligin X Antibody, neuroligin 4 Antibody, X-linked Antibody

Immunogen Fusion protein amino acids 782-945 (intracellular C-terminus) of mouse Neuroligin 4. ~45% identity with Neuroligin-1, -2 and -3.

Purification Protein G Purified

Storage Storage Buffer PBS pH 7.4, 50% glycerol, 0.1% sodium azide

Shipping Temperature Blue Ice or 4ºC Certificate of Analysis

 1μ g/ml of SMC-469 was sufficient for detection of Neuroligin 4 in 20 μ g of mouse brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

-20°C

Cellular Localization Cell Membrane | Cell Junction

Neuroligin 4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

<u>Western Blot</u>

Blocking Peptides



- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Neuroligin 4 Antibody - Images



Immunohistochemistry analysis using Mouse Anti-Neuroligin 4 Monoclonal Antibody, Clone S98-7 (ASM10301). Tissue: Dentate Gyrus. Species: Mouse. Fixation: formaldehyde-fixed paraffin embedded. Primary Antibody: Mouse Anti-Neuroligin 4 Monoclonal Antibody (ASM10301) at 1:100. Secondary Antibody: FITC Goat Anti-Mouse (green). Counterstain: DAPI (blue) nuclear stain at 1:1000. Courtesy of: Rachel Reith, NIH/NIMH.

Neuroligin 4 Antibody - Background

This gene encodes a member of a family of neuronal cell surface proteins. Members of this family may act as splice site-specific ligands for beta-neurexins and may be involved in the formation and remodeling of central nervous system synapses. The encoded protein interacts with discs, large (Drosophila) homolog 4 (DLG4). Mutations in this gene have been associated with autism and Asperger syndrome. Two transcript variants encoding the same protein have been identified for this gene.