

M-CSF/CSF1 Protein, Mouse, Recombinant

#Cat: NB-64-55684-20µg	Size: 20µg
#Cat: NB-64-55684-100µg	Size: 100µg
#Cat: NB-64-55684-200µg	Size: 200µg
#Cat: NB-64-55684-500µg	Size: 500µg

General Information

Synonyms:	Colony stimulating factor 1 (macrophage); C87615;op;Csfm;MCSF
Protein Construction:	A DNA sequence encoding the mouse CSF1 (P07141-1) (Met1-Glu262) was expressed. Predicted N terminal: Lys 33
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	P07141-1
Molecular Weight:	26 kDa (predicted); 45 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	<ol style="list-style-type: none">1. Immobilized mouse CSF1 at 2 µg/ml (100 µl/well) can bind mouse CSF1R-Fch, The EC50 of mouse CSF1R-Fch is 60-220 ng/mL.2. Measured in a cell proliferation assay using M-NFS-60 mouse myelogenous leukemia lymphoblast cells. The ED50 for this effect is typically 3-15 ng/mL.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

Macrophage colony-stimulating factor 1, also known as CSF-1, M-CSF, Lanimostim and CSF1, is a single-pass membrane protein which is disulfide-linked as a homodimer or heterodimer. Granulocyte / macrophage colony stimulating factors are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages. MCSF/CSF-1 is known to facilitate monocyte survival, monocyte-to-macrophage conversion, and macrophage proliferation. M-CSF/CSF-1 is a secreted cytokine which influences hemopoietic stem cells to differentiate into macrophages or other related cell types. It binds to the Colony stimulating factor 1 receptor. M-CSF/CSF-1 may also be involved in development of the placenta. The active form of M-CSF/CSF-1 is found extracellularly as a disulfide-linked homodimer, and is thought to be produced by proteolytic cleavage of membrane-bound precursors. M-CSF/CSF-1 induces cells of the monocyte/macrophage lineage. It also plays a role in immunological defenses, bone metabolism, lipoproteins clearance, fertility and pregnancy. Upregulation of MCSF/CSF-1 in the infarcted myocardium may have an active role in healing not only through its effects on cells of monocyte/macrophage lineage, but also by regulating endothelial cell chemokine expression.

Reference

Pandit J. et al., 1992, Science. 258: 1358-62.

Tokai M. et al., 2000, J Bacteriol. 182 (10): 2865-8.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use