

SARS-CoV-2 N Protein Rabbit mAb, BSA and glycerol free

Catalog No.: A20021 Recombinant 13 Publications

Basic Information

Observed MW

Calculated MW

46kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IF/ICC,ELISA,DB

Cross-Reactivity

SARS-CoV-2

CloneNo number

ARC5077-02

Background

Packages the positive strand viral genome RNA into a helical ribonucleocapsid (RNP and plays a fundamental role during virion assembly through its interactions with the viral genome and membrane protein M. Plays an important role in enhancing the efficiency of subgenomic viral RNA transcription as well as viral replication. May modulate transforming growth factor-beta signaling by binding host SMAD3.

Recommended Dilutions

WB 1:1000 - 1:6000

DB 1:500 - 1:2000

IF/ICC 1:200 - 1:800

ELISA Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

www.abclonal.com

Immunogen Information

Gene ID43740575

Swiss Prot
P59595

Immunogen

Recombinant fusion protein of SARS-CoV-2 Nucleoprotein Full length (P59595) .

Synonyms

Nucleoprotein; NP; SARS-CoV-2 N Protein


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C Avoid freeze / thaw cycles. Buffer: PBS with 0.09% sodium azide,pH7.3.

Validation Data



Western blot analysis of Recombinant SARS-CoV-2 Nucleocapsid Protein (RP01264LQ), using SARS-CoV-2 N Protein Rabbit mAb,BSA and glycerol free (A20021) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 ng/10 ng/5 ng/2 ng per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.