

FAOBlue™ <Fatty Acid Oxidation Detection Reagent>

Catalog NO. FDV-0033

Research use only, not for human or animal therapeutic or diagnostic use.

日本語版はこちらから
ダウンロードできます。
①弊社ウェブサイトより
Webページ番号検索にて
【70873】で検索



Product Background

Fatty acids (FAs) are basic building blocks for wide variety of lipids, essential components of cells, and are one of primary sources of energy. The major pathway for the degradation of FAs is mitochondrial FA beta-oxidation (FAO). FAO is a key metabolic pathway for energy homeostasis in organs such as the liver, heart and skeletal muscle. FAO is a complicated biochemical event containing many types of enzymes. First, FAs are converted to acyl-CoA form by acyl-CoA synthetase family. Second, acyl-CoA forms are incorporated into mitochondria via the carnitine shuttle pathway. Once acyl-CoA entering to mitochondrial matrix, acyl-CoA (C_n) is converted to acyl-CoA (C_{n-2}) and acetyl-CoA by four stepwise reactions. 1) Dehydrogenation: Acyl-CoA is oxidized to enoyl-CoA by acyl-CoA dehydrogenases, 2) Hydration: Enoyl-CoA is hydrated to 3-hydroxyacyl-CoA by crotonase, 3) Oxidation: 3-hydroxyacyl-CoA to 3-ketoacyl-CoA, 4) Thiolysis: 3-ketoacyl-CoA to acyl-CoA (C_{n-2}) and acetyl-CoA. Acetyl-CoA is further converted to ATP. The resulting acyl-CoA (C_{n-2}) enters another cycle of FAO to further produce acyl-CoA (C_{n-4}).

Abnormal FAO is involved in various diseases such as obesity and non-alcoholic fatty liver diseases (NAFLD). Although measurement of FAO activity in diseased cells is important, methods for measuring FAO activity are limited due to its complicated processes above. Only a few indirect methods such as using radio-isotope containing fatty acids or measuring oxygen consumption are commonly performed.




Figure 1. Overview of FAO cycle in mitochondria

FAOBlue™ is the world-first reagent for directly measuring FAO activity in living cells. FAOBlue™ is a coumarin dye possessing a nonanic acid (C9) protected by acetoxymethyl ester, and it shows no fluorescence excited by 405 nm before metabolization by FAO. FAOBlue™ can enter into cells through direct penetration of cell membrane and its acetoxymethyl ester is hydrolyzed by intracellular esterases. Free FA type of FAOBlue™ is converted to acyl-CoA and further incorporated into FAO pathway. Acyl-CoA type of FAOBlue™ is degraded by three FAO cycles to non-fluorescent coumarin possessing a propionic acid (C3). After 4th FAO degradation, coumarin dye is released from propionic acid and diffused into whole cells. While FAOBlue™ shows no fluorescence excited by 405 nm before metabolizing by FAO, the released coumarin derived from FAO cycles shows strong blue fluorescence excited by 405 nm. FAOBlue™ -based FAO assay enables to measure FAO activity with an easy procedure. Inhibition of carnitine shuttle by etomoxir greatly diminishes the fluorescence enhancement in living cells. This data suggests that FAOBlue™ mainly detects mitochondrial FAO activity.




Figure 2. Sensing mechanism of FAOBlue™ in living cell.

Description

Catalog Number: FDV-0033
 Size : 0.2 mg
 Formulation : C₂₄H₃₁NO₉
 Molecular weight : 477.51 g/mol
 Solubility : Soluble in DMSO

Application

- Relative quantification of FAO activity
- Evaluation of drug effect on FAO activity

Reconstitution and Storage

Reconstitution : Add 100% DMSO into vial to prepare 1-10 mM stock solution.

Storage :

Powder : Store at -20°C.


Stock solution :

- Make aliquots and store at -20 °C with protecting from light.
- Avoid repeated freeze-thaw cycles.

How to use

General procedure

1. Add FAOBlue™ (recommended final conc. 5-20 μM) in fresh HEPES-buffered saline (HBS; cf. 20 mM HEPES (pH 7.4), 107 mM NaCl, 6 mM KCl, 1.2 mM MgSO_4 , 2 mM CaCl_2)
NOTE: Instead of HBS, serum free-/phenol red free-culture media are also suitable. As serum proteins may absorb FAOBlue™ and inhibit cellular uptake of FAOBlue™, serum free is recommended. In cases of using serum free culture media, as nutrients in culture media such as glucose, pyruvate etc. may affect cellular FAO activity, please empirically optimize reagent conc., incubation time etc. for each experiment.
2. Remove the cultured medium and wash the cells with HBS (or serum free culture media) twice
3. Add FAOBlue™ -containing HBS (or serum free culture media) to cells
4. Incubate cells at 37°C for >30 min
5. Wash cells with HBS (or culture media)
6. Observe cells under live condition with blue fluorescence (Ex 405 nm/ Em. 430-480 nm)



Reference data and Experimental guide


Absorbance and fluorescence spectrum of FAOBlue™ and coumarin-derivative

Absorbance (left) and fluorescence (right) spectra of FAOBlue™ (blue line) and the coumarin-derivative dye released after metabolization of FAO (red line) in PBS buffer (pH 7.4).

Absorbance spectra: an absorption peak of coumarin-derivative dye is clearly shifted by FAO from the peak of FAOBlue™.

Fluorescence spectra: Coumarin-derivative dye shows strong blue fluorescence but FAOBlue™ emits little fluorescence, when both compounds are excited at 405 nm.

NOTE: FAOBlue™ shows blue fluorescence (gray line; 370-450 nm) when it is excited at 300-380 nm (max 350 nm).



Experimental guide for fluorescence imaging

Confocal laser microscopy: Please use 405 nm laser equipped in the microscopy. Using 405 nm laser allows to detect only a fluorescent signal from coumarin-derivative dye.

Epifluorescence microscopy: Excitation filter is very important. Commercial DAPI filters are not compatible with this reagent, because DAPI filters excite both FAOBlue™ and coumarin-derivative dye. Excitation filters which pass 390-450 nm wavelength light are recommended.

NOTE: Around 400 nm excitation may induce autofluorescence from cells. Negative control (without FAOBlue™) for each experiment is highly recommended to confirm cell-derived autofluorescence signals.


Application data

Visualization of FAO activities in 4 cancer cell lines

Four cancer cell lines (HepG2, LNCaP, HeLa and A549) were treated with FAOBlue™ in HBS buffer with or without pre-treatment of etomoxir (40 μM, 3 hours), a potent FAO inhibitor. After FAOBlue™ incubation, blue fluorescence (Ex. 405 nm/ Em. 430-480 nm) was observed. All cell lines showed blue fluorescence in cytosol, but pre-treatment of etomoxir clearly decreased fluorescent intensities. These results indicated the blue fluorescence was derived from FAO activity in the cells.


* Experimental condition:

- HepG2, 5 μM FAOBlue™ for 30 min.
- LNCaP, 20 μM FAOBlue™ for 120 min
- HeLa, 20 μM FAOBlue™ for 120 min
- A549, 5 μM FAOBlue™ for 30 min




Perturbation of FAO activity by drugs

HepG2 cells were pre-incubated with 200 μM AICAR, a FAO activator via AMPK activation, for 3 hours or 200 μM of ranolazine, a partial FAO inhibitor, for 12 hours. After drug treatment, the cells were incubated with 5 μM FAOBlue™ for 30 min. Compared with control cells, pretreatment with AICAR significantly increased blue fluorescent intensity. On the other hand, pretreatment with the partial FAO inhibitor ranolazine clearly decreased blue fluorescent intensity. Ex. 405 nm/ Em. 430-480 nm



Quantitative analysis of the drug effects on FAO activity


ND630 is an inhibitor of acetyl-CoA carboxylase and considered as a potential therapeutic drug for non-alcoholic fatty liver disease (NAFLD). HepG2 cells were pre-incubated with various concentration of ND630 for 4 hours. After ND630 treatment, cells were treated with 5 μ M FAOBlue™ for 30 min. Blue fluorescent intensities of each concentration of ND630 were quantified. Ex. 405 nm/ Em. 430-480 nm



Analysis of FAO activity using NASH model mouse

Non-alcoholic steatohepatitis (NASH) is a typical disease which shows low metabolic activity of FAs. Control healthy mice and NASH model mice were orally administered with 400 mg/kg bezafibrate, a therapeutic agent of NASH. After 4 weeks administration, primary hepatocytes were isolated from control mice and NASH model mice and cultured in culture dishes. Primary hepatocytes were further treated with 5 μ M FAOBlue™ for 30 min and fluorescence imaging was performed. Compared with control cells, NASH model mouse-derived hepatocytes showed low FAO activity. Bezafibrate dramatically recovered FAO activity of hepatocytes isolated from NASH model mouse. FAOBlue™ is a powerful tool to estimate drug effects and efficiency on FAO activity.


*Detail procedure of mice experiment is described in Ref.1. Ex. 405 nm/ Em. 430-480 nm



High-throughput cell-based detection by fluorescent plate reader

786-O cells were seeded in 96 well and culture for 1 day. Confluent cells were treated with indicated drugs in serum- and phenol red-free DMEM for 20 hours. After wash cells with HBS, cells were treated with final 10 μM of FAOBlue™ in HBS for 2 hours. Without washing cells, fluorescent intensity (Ex 420 \pm 5 / Em 460 \pm 10) by the fluorescent plate reader with a transparent mode. Fluorescent signals were normalized by the background signal of 10 μM FAOBlue™ without cells. A mitochondrial FAO inhibitor, etomoxir, clearly suppress blue fluorescent signal.

[Drug treatment: 10 μM Etomoxir (mitochondrial FAO inhibitor), 500 μM AICAR (AMPK activator), 5 μM 2-bromopalmitate (2-BP; broad lipid metabolism inhibitor)]



Reference

1. Uchinomiya *et al.*, *Chem. Commun.*, **56**, 3023-3026 (2020) Fluorescence Detection of Metabolic Activity of Fatty Acid Beta Oxidation Activity in Living Cells.

Disclaimer/免責事項

This product has been commercialized by Funakoshi Co., Ltd. based on the results of academic research, and the advertisement text, figures and manuals (hereinafter “Product information”) have been prepared based on published research reports on February, 2020. The academic interpretation at the time of creation of the Product Information may change in accordance with future developments in the relevant research field and expansion of various scientific findings, and the latest version and certainty of the Product Information are not guaranteed. The specifications of this product and the Product Information are subject to change without notice. Please contact us for the latest information.

本商品は学術研究成果を基にフナコシ株式会社が商品化したもので、2020年2月時点における公開研究報告を基に広告文章およびマニュアル(以下、商品資料)を作成しています。今後の当該研究分野の発展および各種学術知見の拡大にともない、商品資料作成時の学術的解釈が変更になる可能性があり、最新性・確実性を保証するものではありません。また、本商品の仕様および商品資料を予告なく変更する場合がございます。最新の情報に関しましては、弊社までご確認いただけますようお願い申し上げます。

E-mail Newsletter Sign Up

Japanese English

Related products

Lipidye™ II <Lipid Droplet Live Imaging>


Lipidye™ II is a highly sensitive lipid droplet staining dye with extremely photostable property. This dye is the second generation of our previous reagent, Lipidye™. This dye allows us to detect small lipid droplets (<1 μm) in non-adipocytes and to apply into long-term live cell imaging for dynamic lipid droplet movements.

Catalog No. FDV-0027

Size 0.1 mg

Features

- Recommended Ex/Em:400-500 nm / 490-550 nm
- Enable to detect <1 μm lipid droplets
- Suitable for long-term live cell imaging
- Extremely photostable compared with conventional dyes
- Compatible with both live and fixed cells



Lipidye™-M <Lipid Metabolism Tracer>


Lipidye™-M is a C12 fatty acid mimic labeled with a novel solvatochromic dye. As Lipidye™-M exhibits green-to-red fluorescence depending on its lipid structure and its localization, Lipidye™-M can trace status of cellular fatty acid uptake and lipid metabolism in cells. Lipidye™-M is a powerful tool for both basic research and pharmaceutical research for lipid metabolism.

Catalog No. FDV-0028

Size 0.1 mg

Features

- allows to perform three-color imaging (green, yellow and red) by merging images from a green channel (Ex. 450-490 nm / Em. 490-540 nm) and red channel (Ex. 550-600 nm / Em. 570-620 nm)
- can also be taken up to cells by FA-transporters and converted into many types of lipids, including acyl-CoA, phospholipids, DAGs, TAGs, and degraded to small metabolites by the mitochondrial FAO pathway.
- Emits green fluorescence in high polarity (cytosol), yellow fluorescence in moderate polarity (organelle membranes) and red fluorescence in low polarity (lipid droplets)



LipiRADICAL™ Green <Lipid Radical Detection Reagent>


LipiRADICAL™ Green is a specific fluorescent dye for lipid-derived radicals which are the most upstream factor of lipid peroxidation (LPO). LipiRADICAL™ Green can be applied into both *in vitro* assay and cell-based assay to monitor lipid radical productions.

Catalog No. FDV-0042

Size 0.1 mg

Features

- Recommended Ex/Em: ~480 nm / 520 nm
- Enable to detect very unstable lipid-derived radicals
- Compatible with *in vitro* assay and in cell-based assay
- An innovative reagent for comprehensive identification of lipid-derived radicals by lipidomics



LipiORDER™ <Membrane Lipid Order Imaging Dye>

LipiORDER™ is a solvatochromic dye for membrane lipid order imaging. LipiORDER™ exhibits green fluorescence with Lo phase and exhibits red fluorescence with Ld phase. The ratiometric analysis (F_{red}/F_{green}) enables the quantitative visualization of membrane lipid order.

Catalog No. FDV-0041

Size 0.1 mg

Features

- Recommended Ex/Em: ~405 nm / 500-550 nm (Green channel) and 550-650 nm (Red channel)
- Enable to quantitatively monitor lipid order on plasma and inner membranes in live cells
- Highly photostable and cellularly stable compared with similar conventional dyes.

