

RP02324

Leader in Biomolecular Solutions for Life Science



Recombinant SARS-COV-2 Spike S1 (N501Y) Protein

Catalog No.: RP02324 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	43740568	YP_009724390.1

Tags

C-His

Synonyms

S1 protein;Spike glycoprotein
Subunit1;S glycoprotein
Subunit1;Spike protein S1

Product Information

Source	Purification
HEK293 cells	> 95% by Tris-Bis PAGE;> 95% by SEC-HPLC

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

Basic Information

Description

Recombinant SARS-COV-2 SARS-COV-2 Spike S1 (N501Y) Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Val16-Arg685) of SARS-COV-2 SARS-COV-2 Spike S1 (N501Y) fused with His tag at the C-terminal.


Bio-Activity

Immobilized SARS-COV-2 Spike S1 (N501Y) ,His Tag at 0.5 μg/mL (100 μL/Well) on the plate. Dose response curve for Human ACE2 ,hFc Tag with the EC₅₀ of 80.3 ng/mL determined by ELISA.


Storage

Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Validation Data



SARS-COV-2 Spike S1 (N501Y) on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of SARS-COV-2 Spike S1 (N501Y) is greater than 95% as determined by SEC-HPLC.



Immobilized SARS-COV-2 Spike S1 (N501Y), His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Human ACE2, hFc Tag with the EC₅₀ of 80.3ng/ml determined by ELISA.