

For research use only

KO616 Anti mouse AIM Monoclonal Antibody

Clone No. 20C1

Target mouse AIM
Category immunology
Gene ID 11801

Primary Source MGI:1334419

Synonyms CD5L, AAC-11, AIM/Spalpha, Api6, Pdp 1/6, Sp-alpha

Type Monoclonal Antibody
Immunogen recombinant mouse AIM

Raised in Wistar Rat

Myeloma P3U1

Clone number 20C1 (#29)

Purification ProteinG

Source Serum-free medium

 $\begin{tabular}{lll} Isotype & IgG1\kappa \\ \hline Cross Reactivity & Not tested \\ \hline Label & Unlabeled \\ \hline \end{tabular}$

Concentration Contents(Volume) Buffer PBS

Storage Store at - 20°C long term, store at 4°C short term. Avoid repeated freeze-thaw cycles.

Application ELISA, WB, ICC, IP

ELISA	WB	IHC	ICC
1.0	1.0	Not tested	1.0
IP	FCM	IF	Neutralization
5.0	Not tested	Not tested	ı

(µg/mL)

Reference

Miyazaki T et al. AlMing at Metabolic Syndrome—Towards the Development of Novel Therapies for Metabolic Diseases via Apoptosis Inhibitor of Macrophage (AIM) –Circ. J., 2011, 75, 2522-2531

Kurokawa et al. Apoptosis inhibitor of macrophage (AIM) is required for obesity-associated recruitment of inflammatory macrophages into adipose tissue. Proc Natl Acad Sci USA 2011, 108, 12072-12077

Kurokawa et al. Macrophage-derived AIM is endocytosed into adipocytes and decreases lipid droplets via inhibition of fatty acid synthase activity. Cell Metab. 2010, 11, 479-492

UniProt Summary

//Function: May play a role in the regulation of the immune system. Seems to play a role as an inhibitor of apoptosis.

//Subcellular location:Secreted.

//Tissue specificity: Expressed in thymus, liver, spleen and lymph nodes.

//Post-translational modification: Glycosylated.

//Sequence similarities: Contains 3 SRCR domains.